

Sudden death of distillability in qutrit-qutrit systems

Wei Song,^{1,*} Lin Chen,^{2,†} and Shi-Liang Zhu¹

¹*Laboratory of Quantum Information Technology, ICMP and SPTE,
South China Normal University, Guangzhou 510006, China*

²*Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117542*
(Dated: 2 Jul 2009)

We introduce the concept of distillability sudden death, i.e., free entangled states can evolve into non-distillable (bound entangled or separable) states in finite time under local noise. We describe the phenomenon through a specific model of local dephasing noise and compare the behavior of states in terms of the Bures fidelity. Then we propose a few methods to avoid distillability sudden death of states under (general) local dephasing noise, so that free entangled states can be robust against decoherence. Moreover, we find that bound entangled states are unstable in the limit of infinite time.

PACS numbers: 03.65.Yz, 03.67.Mn, 03.65.Ud, 03.67.Pp

I. INTRODUCTION

Entanglement is not only a remarkable feature which distinguishes the quantum world from the classical one but also a key resource to realize high-speed quantum computation and high-security quantum communication [1]. In realistic quantum-information processing, entanglement usually needs to be prepared or distributed beforehand among different remote locations. However, in the process of entanglement distribution the quantum systems are not isolated and each system will unavoidably interact with the environment. This leads to local decoherence which will degrade the entanglement of the shared states. It is thus of fundamental importance to study the entanglement properties under the influence of the local decoherence. In this context, Yu and Eberly [2] investigated the time evolution of entanglement of a bipartite qubit system undergoing various modes of decoherence. Remarkably, they found that, although it takes infinite time to complete the decoherence locally, the global entanglement may vanish in finite time. The phenomenon of finite-time disentanglement, also named entanglement sudden death (ESD), unveils a fundamental difference between the global behavior of an entangled system and the local behavior of its constituents under the effect of local decoherence. Clearly, ESD puts a limit on the applicability of entangled states in the practical quantum information processing.

Initially, Yu and Eberly reported the ESD for two-qubit entangled states, but this effect is not limited to such case. Further investigations in a wider context including higher dimensional Hilbert spaces have been made by various groups [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]. There are also a number of studies looking at ESD in more complicated systems using other entanglement measures [20, 21, 22, 23, 24, 25, 26, 27],

and an attempt to give a geometric interpretation of the phenomenon has also been made [28]. In addition, experimental evidences of ESD have been reported for optical setups [29] and atomic ensembles [30].

However, all previous studies omitted an important fact that high dimensional bipartite entangled states can be divided into two classes [31]. One is free, which means that the state can be distilled under local operations and classical communication (LOCC); the other is bound, which means that no LOCC strategy is able to extract pure-state entanglement from the state even if many copies are available. Bound entanglement (BE) cannot be used alone for quantum information processing, and irreversibility occurs in asymptotic manipulations of entanglement for all BE states [32]. Since it was constructed from a pure mathematical point of view, we may ask whether BE can appear in physically relevant quantum systems naturally. Very recently a few works have addressed this question [33]. Their results suggest that different many-body models present thermal bound entangled states.

In this paper, we investigate the problem from a very different viewpoint, i.e., in the presence of local decoherence, which is one of the dominant noises during the distribution of entanglement. Analogous to the definition of ESD, if an initial free entangled state becomes non-distillable in finite time under the influence of local decoherence, then we say that it undergoes distillability sudden death (DSD). Note that when a free entangled state loses its distillability at a specific time, it may still be entangled since we cannot exclude the existence of BE. So far, it is not clear whether bipartite bound entangled states can be created from free entangled states under local decoherence process naturally. The first aim of this paper is to show that such a process indeed exists through an explicit qutrit-qutrit example.

Afterwards we propose the DSD-free state, which has entanglement robust against local decoherence. Such entangled states are thus useful resources for practical quantum-information processing. So the second aim is to address the DSD-free state. We develop a few sys-

*Electronic address: wsong1@mail.ustc.edu.cn

†Electronic address: cqtc1@nus.edu.sg (Corresponding Author)

tematic approaches to build DSD-free states. Finally we will show that different from free entangled and separable states, no PPT bound entangled states exist in the infinite time limit.

The paper is organized as follows. In Sec. II, the idea of DSD is presented. In particular, we show the phenomenon of DSD through a specific qutrit-qutrit example. In addition, we show that one can avoid the sudden death of distillability by performing a simple local unitary operation on the initial state. Furthermore, we compare the behaviors of states affected by decoherence in terms of the Bures fidelity. In Sec. III, we develop some methods to build DSD-free states which can protect free entanglement under general local dephasing noise. We also show that no PPT entangled states can exist in the limit of infinite time. Finally, in Sec. IV we discuss some open questions and also give a summary of our results.

II. QUTRIT-QUTRIT DSD STATES UNDER LOCAL DECOHERENCE

Before discussing dynamical process of entanglement, we briefly review how to characterize bound entangled states. It was proven in [31] that a quantum state with a positive partial transposition (PPT) is non-distillable under LOCC. Therefore PPT entangled states must be bound entangled states [34]. To verify them, one can use the so-called realignment criterion (cross-norm criterion) [35]. The definition of realignment on the density matrix is given by $(\rho^R)_{ij,kl} = \rho_{ik,jl}$. A separable state ρ always satisfies $\|\rho^R\| \leq 1$. For a PPT state ρ , the positive value of the quantity $\|\rho^R\| - 1$ can hence verify that it is a bound entangled state.

The system we study consists of two noninteracting qutrits in two independent local environments, each coupling to one of the qutrits. Here the qutrit is a three-dimensional state composed of the computational basis $|0\rangle$, $|1\rangle$, and $|2\rangle$. Besides, we take the weak local dephasing noise to model the environment, such noise is indeed one of the main decoherence sources in solid-state systems.

The general time-evolved density matrix expressible in the operator-sum decomposition is the completely positive trace preserving map [36] $\rho(t) = \mathcal{E}(\rho(0)) = \sum_{\mu} K_{\mu}^{\dagger}(t) \rho(0) K_{\mu}(t)$. The operators $\{K_{\mu}(t)\}$ representing the influence of statistical noise satisfy the completeness condition $\sum_{\mu} K_{\mu}(t) K_{\mu}^{\dagger}(t) = \mathbb{I}$ which guarantees that the evolution is trace-preserving[1]. In our model, the operators are of the form $K_{\mu}(t) = D_j(t) E_i(t)$ such that

$$\rho(t) = \mathcal{E}(\rho(0)) = \sum_{i,j=1}^2 D_j^{\dagger}(t) E_i^{\dagger}(t) \rho(0) E_i(t) D_j(t). \quad (1)$$

Here, $E_i(t)$ and $D_j(t)$ correspond to local dephasing noise components acting on the first and second qutrits, respectively, and both operators satisfy the completeness




FIG. 1: (Color online). The eigenvalue λ_1 as a function of time t and dephasing rate Γ for two different cases: (a) $\alpha = 4.5$; (b) $\alpha = 4.9$. Figures (c) and (d) correspond to the contour plots of the upper one, respectively.

conditions. For simplicity, we first take these to be of the specific forms [14]

$$\begin{aligned} E_1(t) &= \text{diag}(1, \gamma_A, \gamma_A) \otimes \mathbb{I}_3, \\ E_2(t) &= \text{diag}(0, \omega_A, \omega_A) \otimes \mathbb{I}_3, \\ D_1(t) &= \mathbb{I}_3 \otimes \text{diag}(1, \gamma_B, \gamma_B), \\ D_2(t) &= \mathbb{I}_3 \otimes \text{diag}(0, \omega_B, \omega_B), \end{aligned}$$

where $\mathbb{I}_3$ is the 3×3 identity matrix, $\gamma_A(t) = e^{-\Gamma_A t/2}$, $\gamma_B(t) = e^{-\Gamma_B t/2}$, $\omega_A(t) = \sqrt{1 - \gamma_A^2(t)}$, and $\omega_B(t) = \sqrt{1 - \gamma_B^2(t)}$.

For concreteness, we illustrate our ideas by considering the following qutrit-qutrit state,

$$\begin{aligned} \rho(0) &= \frac{2}{21} (|01\rangle + |10\rangle + |22\rangle) (\langle 01| + \langle 10| + \langle 22|) \\ &+ \frac{\alpha}{21} (|00\rangle \langle 00| + |12\rangle \langle 12| + |21\rangle \langle 21|) \\ &+ \frac{5-\alpha}{21} (|11\rangle \langle 11| + |20\rangle \langle 20| + |02\rangle \langle 02|) \end{aligned} \quad (2)$$

with $4 < \alpha \leq 5$. In fact, it is straightforward to prove that the state $\rho(0)$ is free entangled. By using the local operation $(|0\rangle \langle 0| + |1\rangle \langle 1|) \otimes (|0\rangle \langle 0| + |1\rangle \langle 1|)$, one can convert $\rho(0)$ into a $2 \otimes 2$ entangled state, which is afterwards distillable by virtue of the BBPSSW-Horodecki protocol [37]. So the state $\rho(0)$ is a free entangled state and we take it as the initial state under the local dephasing noise in Eq. (1). This evolution can be calculated analytically, i.e., at time t the two-qutrit density operator $\rho(t)$ reads

$$\begin{pmatrix} \frac{\alpha}{21} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{21} & 0 & \frac{2}{21}\gamma_A\gamma_B & 0 & 0 & 0 & 0 & \frac{2}{21}\gamma_A \\ 0 & 0 & \frac{5-\alpha}{21} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2}{21}\gamma_A\gamma_B & 0 & \frac{2}{21} & 0 & 0 & 0 & 0 & \frac{2}{21}\gamma_B \\ 0 & 0 & 0 & 0 & \frac{5-\alpha}{21} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{\alpha}{21} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{5-\alpha}{21} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{\alpha}{21} & 0 \\ 0 & \frac{2}{21}\gamma_A & 0 & \frac{2}{21}\gamma_B & 0 & 0 & 0 & 0 & \frac{2}{21} \end{pmatrix}, \quad (3)$$

where the basis are spanned by $\{|00\rangle, |01\rangle, |02\rangle, |10\rangle, |11\rangle, |12\rangle, |20\rangle, |21\rangle, |22\rangle\}$.

Let us analyze the matrix carefully. There are three eigenvalues of the partial transposition of the state $\rho(t)$ which could be negative, namely $\lambda_1 = f(\Gamma_A)$, $\lambda_2 = f(\Gamma_B)$, and $\lambda_3 = f(\Gamma_A + \Gamma_B)$, where $f(\lambda) = \frac{e^{-\lambda t}}{882} \left(105e^{\lambda t} - \sqrt{11025e^{2\lambda t} + 1764e^{\lambda t}(4 - 5\alpha e^{\lambda t} + \alpha^2 e^{2\lambda t})} \right)$.

For simplicity, we choose the local asymptotic dephasing rates $\Gamma_A = \Gamma_B = \Gamma$, and thus $\lambda_1 = \lambda_2 < \lambda_3$ in the following arguments. In order to have a vivid illustration, in Fig. 1 we plot λ_1 as a function of t and Γ with specific $\alpha = 4.5$ and $\alpha = 4.9$, respectively. Fig. 2(a) shows the value of λ_1 versus t for different decoherence rates with $\alpha = 4.5$, and we can see that the eigenvalue of the partial transposition of the state $\rho(t)$ will always arrive at a positive value in finite time. For example, if we choose $\Gamma = 1$, the density matrix $\rho(t)$ will become a PPT state after time $t \approx 0.58$ in Fig. 2(a). Analytically, the time at which $\rho(t)$ becomes a PPT state is $t_d = \frac{1}{\Gamma} \ln \frac{4}{\alpha(5-\alpha)}$.

Next, we use realignment to verify the BE in this evolution. To this end, we need to compute the quantity $\|\rho(t)^R\| - 1$, and it is given by $\frac{2}{21}e^{-\Gamma t} \left(2 + 4e^{\frac{1}{2}\Gamma t} + (-7 + \sqrt{19 - 15\alpha + 3\alpha^2})e^{\Gamma t} \right)$. In order to make a comparison, we plot $\|\rho(t)^R\| - 1$ as a function of time t by choosing $\alpha = 4.5$ and $\Gamma = 1$ in Fig. 2(b). We can see that if $t < 0.84$, the value of $\|\rho(t)^R\| - 1$ is always positive, which indicates that in the range $0.58 < t < 0.84$, the two-qutrit system is a bound entanglement state. Thus we have shown that free initial entangled states can evolve into non-distillable (bound) entangled states in finite time under the local external asymptotic dephasing noise.

Besides we find that with $\alpha = 4.5$ and $\Gamma = 1$, the state $\rho(t)$ will be separable when $t > 1.39$. It could be proved by extracting three 2×2 density operators, which are respectively spanned by $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$, $\{|01\rangle, |02\rangle, |21\rangle, |22\rangle\}$ and $\{|10\rangle, |12\rangle, |20\rangle, |22\rangle\}$, from the state $\rho(t)$. Then one can find out that the condition $t > 1.39$ makes the three states separable. As the state $\rho(t)$ is actually the linear combination of the above three states and some product states, $\rho(t)$ also becomes separable when $t > 1.39$. Nevertheless, there is still a small window $t \in [0.84, 1.39]$ in which the state $\rho(t)$ has merely PPT and we don't know whether it's entangled




FIG. 2: (a) The eigenvalue λ_1 versus time t for different dephasing rates. The solid, dash-dotted, and dashed lines correspond to $\Gamma = 1$, $\Gamma = 0.7$, and $\Gamma = 0.4$, respectively. (b) The quantity $\|\rho(t)^R\| - 1$ versus time t for $\Gamma = 1$. The other parameter $\alpha = 4.5$.

or separable.

To summarize the example, start from the initial state $\rho(0)$ in $4 < \alpha \leq 5$ the qutrit-qutrit state $\rho(t)$ will experience three phases under the local dephasing noise in Eq.(1): it evolves from an initial distillable entangled state to a PPT entangled state, and then changes to be a separable state and finally remains in this phase for even. In other word, $\rho(t)$ first experiences DSD and then ESD under local decoherence.

In contrast with the above example, we briefly consider the initial state $\rho(0)$ for $3 < \alpha \leq 4$. In this case, it is a PPT entangled state [38], one can verify that the state $\rho(t)$ will finally become separable in finite time under the local dephasing noise in Eq.(1); that is, $\rho(t)$ only experiences ESD. Moreover, we can easily verify the time-domain factorization relation as follows,

$$S_{t_1+t_2} = S_{t_1}S_{t_2}, \quad (4)$$

where $t_1, t_2 \geq 0$, and S denotes the noise in Eq.(1). Because the entanglement cannot increase under the local operations, we conclude that if a state becomes separable at a specific time in Eq.(1), it must remain separable in all subsequent time. More generally, if an entangled state evolves into a PPT state (either entangled or separable) at a specific time in Eq.(1), it always has PPT for all subsequent time since the local operation cannot change PPT.

Of course, such disappearance of distillability in finite time can seriously affect the application of entanglement in quantum information tasks. An important question arises naturally: given a free initial entangled state, does there exist a suitable intervention that may alter the final fate of DSD? In the following we show that one can realize this aim by merely performing a simple local unitary operation on the initial state in the above example. Let the initial local operation be $U = \mathbb{I}_3 \otimes \lambda$, with $\lambda = |0\rangle\langle 1| + |1\rangle\langle 0| + |2\rangle\langle 2|$.

Then, the transformed state is $\rho'(0) = U\rho(0)U^\dagger = \frac{2}{7}P_+ + \frac{\alpha}{7}\rho_+ + \frac{5-\alpha}{7}\rho_-$, where the projectors are $P_+ = |\Psi_+\rangle\langle\Psi_+|, |\Psi_+\rangle = \frac{1}{\sqrt{3}}(|00\rangle + |11\rangle + |22\rangle), \rho_+ = \frac{1}{3}(|01\rangle\langle 01| + |12\rangle\langle 12| + |20\rangle\langle 20|), \rho_- = \frac{1}{3}(|10\rangle\langle 10| + |21\rangle\langle 21| + |02\rangle\langle 02|)$.

Evidently, the states $\rho'(0)$ and $\rho(0)$ are equally useful quantum resources without decoherence. In Ref.[38], Horodecki demonstrated that $\rho'(0)$ is a free entangled state for $4 < \alpha \leq 5$. It will evolve into $\rho'(t)$ according to Eq.(1), whose partial transpose always has a negative eigenvalue in finite time t . So $\rho'(t)$ never experiences ESD when subjecting only to the local dephasing noise in Eq.(1). Furthermore, the state $\rho'(t)$ does not undergo DSD in finite time. To see this, we perform the local operation $(|1\rangle\langle 1| + |2\rangle\langle 2|) \otimes (|1\rangle\langle 1| + |2\rangle\langle 2|)$ on $\rho'(t)$. The resulting two-qubit state is easily proved to be entangled and hence distillable in finite time t . So the state $\rho'(t)$ can always be distilled under the local dephasing noise.

By far we have considered DSD in the presence of local dephase noise. Another useful physical quantity in quantum-information problems is the fidelity, which measures to what extent the evolved state is close to the initial one. For concreteness, we study the states $\rho(t)$ and $\rho'(t)$ by means of the Bures fidelity [39]. The Bures fidelity of states ρ and σ is defined as $F(\rho, \sigma) = [\text{tr}(\sqrt{\sqrt{\rho}\sigma\sqrt{\rho}})]^2$. For the state $\rho(t)$, the fidelity is given by

$$F^\rho(t) = \left[\frac{1}{21} \left(15 + \sqrt{6e^{-\Gamma t} (1 + 2e^{\Gamma t} + \sqrt{1 + 8e^{\Gamma t}})} \right) \right]^2, \quad (5)$$

while for $\rho'(t)$ it reads

$$F^{\rho'}(t) = \left[\frac{1}{21} \left(15 + \sqrt{18 + 6\sqrt{1 + 8e^{-2\Gamma t}}} \right) \right]^2. \quad (6)$$

Eq.(5) and Eq.(6) indicate that the fidelities do not depend on the coefficient α . In Fig.3, we plot $F^\rho(t)$ and $F^{\rho'}(t)$ as the function of t with $\Gamma = 1$. Obviously, the fidelity $F^{\rho'}(t)$ is always larger than $F^\rho(t)$, which means that $\rho(t)$ degrades faster than $\rho'(t)$. In the infinite time limit, the values of $F^{\rho'}(t)$ and $F^\rho(t)$ will approach to $[\frac{1}{21}(15 + 2\sqrt{6})]^2$ and $[\frac{1}{21}(15 + 2\sqrt{3})]^2$, respectively. Thus, we have shown that $\rho'(t)$ have the advantages over $\rho(t)$ in two aspects: Firstly, it does not experience DSD in finite time. Secondly, its fidelity decays slower than that of $\rho(t)$ for all time.

III. CONSTRUCTION OF DSD-FREE STATES

In the last section we have presented the examples of qutrit-qutrit states with and without DSD. We call the latter as DSD-free states, which always have negative




FIG. 3: The fidelity versus time t for different initial states. The solid and dashed lines correspond to $F^\rho(t)$ and $F^{\rho'}(t)$ with $\Gamma = 1$, respectively.

partial transposition (NPT). NPT is a necessary condition for distillability under LOCC, so the DSD-free states are available resource for quantum-information tasks as they have robust entanglement against local dephasing noise described in Eq.(1). It is thus worth finding out diverse DSD-free states in terms of different kinds of local dephasing noises. In this sense, DSD-free states deserve storage in quantum-information "warehouse" and they are more useful than DSD states. On the other hand, one can also directly propose schemes of quantum information processing based on the robustness of DSD-free states.

We now turn to develop several approaches to construct DSD-free states for an arbitrary qutrit-qutrit state σ . As a direct method, we can locally project a few copies of the state $\sigma(t)$ onto some $2 \otimes N (N \geq 2)$ states. Such states are distillable (DSD-free) under LOCC, if and only if they have NPT in finite time [37]. Generally, it's difficult to find out the projectors and there have been a few results on this problem [34, 40, 41]. Here we only investigate one copy of $\sigma(t)$ under the dephasing noise described in Eq.(1). For example:

Lemma 1. Consider an NPT entangled state $\sigma(0)$ and it experiences the local dephasing noise in Eq.(1). Then $\sigma(t)$ is always distillable in finite time, when $\sum_{i=1}^2 D_1^\dagger(t) E_i^\dagger(t) \sigma(0) E_i(t) D_2(t)$ or $\sum_{i=1}^2 D_i^\dagger(t) E_2^\dagger(t) \sigma(0) E_2(t) D_i(t)$ is entangled.

Proof. One can get the above given state by projecting $\sigma(t)$ with the projector $D_2(t)$ on system A or $E_2(t)$ on system B respectively (both projectors actually have nothing to do with time t when separately used as operators). Since the given state is a distillable qubit-qutrit entangled state, the assertion follows from [37]. It's interesting to note that the given state is just the sum of the third and fourth terms in Eq.(1). ■

Lemma 1 actually requires to check the PPT of some 6×6 matrices depending on time t , which could be done by using Mathematics. This is a systematic way to build the subspace of robust entangled states that never experience DSD in Eq. (1). To show the connection between the last section and lemma 1, we propose a weak version of lemma 1 as follows.

Lemma 2. Suppose $D_2^\dagger(t) E_2^\dagger(t) \sigma(0) E_2(t) D_2(t)$ are

entangled, then the states $\sigma(0)$ are DSD-free states in local dephasing noise Eq.(1).

Proof. The state $\sum_{i=1}^2 D_2^\dagger(t) E_i^\dagger(t) \sigma(0) E_i(t) D_2(t)$ is entangled when $D_2^\dagger(t) E_2^\dagger(t) \sigma(0) E_2(t) D_2(t)$ is entangled. The assertion then follows from lemma 1. ■

One may see that the state $\rho'(0)$ in the previous section belongs to the case of lemma 2. The advantage of lemma 2 is that the density operators $D_2^\dagger(t) E_2^\dagger(t) \sigma(0) E_2(t) D_2(t)$ do not contain time t and are convenient for calculation. This result is reasonable for the local dephasing noise described in Eq.(1) which occurs only between the ground state $i = 0$ and the i th ($i = 1, 2$) excited state.

The model in Eq.(1) is neither the simplest case of local dephasing nor the most general case. In the most general case dephasing occurs between all local basis states within each subsystem. In what follows we put forward some results under such general local dephasing noise [14].

Lemma 3. Qutrit-qutrit entangled MC states are DSD-free states in general local dephasing noise.

Proof. Let us consider the qutrit-qutrit maximally correlated (MC) states $\sigma_{MC}(0)$, which read [42]

$$\sigma_{MC}(0) = \sum_{i,j=0}^2 a_{ij} |ii\rangle\langle jj|. \quad (7)$$

We make it go through the channel of general local dephasing noise. As we know, such noise cannot change the diagonal elements and the non-diagonal elements of the evolved state have non-zero dephasing values for any finite time. So the evolved state $\sigma_{MC}(t)$ is still a MC state. On the other hand, the MC state is separable if and only if all non-diagonal entries are zero. Then $\sigma_{MC}(t)$ is always entangled for all finite t whenever the initial state $\sigma_{MC}(0)$ is entangled. Furthermore, one may prove that $\sigma_{MC}(t)$ can always be distillable by using the similar argument to show the distillability of the state $\rho'(t)$ in the previous section. ■

This example enlightens us on building DSD-free entangled states in a stronger way, i.e.,

Lemma 4. We perform the local operation $(|i\rangle\langle i| + |j\rangle\langle j|) \otimes (|m\rangle\langle m| + |n\rangle\langle n|)$ on any initial qutrit-qutrit state, where i, j, m , and n are chosen from the basis 0, 1 and 2, respectively. If the projected state is an entangled MC state, then the initial state is a DSD-free state in general local dephasing noise. ■

One can prove lemma 4 by showing that there are always some nonzero non-diagonal entries of the projected state, and we can take it as the new initial state in the local dephasing noise. Evidently, the newly built state in lemma 4 contains MC states in lemma 3.

These results can be extended to the scenario of higher dimension. For simplicity, we consider the general MC state $\sigma'_{MC}(0) = \sum_{i,j=0}^{d-1} a_{ij} |ii\rangle\langle jj|$. Suppose it experiences generalized local dephasing noise. Analogous to the arguments for lemma 3, one can see that the state $\sigma'_{MC}(t)$

also has no DSD in finite time. Thus we have provided a high dimensional subspace $\{|00\rangle, |11\rangle, \dots, |d-1, d-1\rangle\}$ in which all entangled states are distillable under local dephasing noise; that is,

Lemma 5. Entangled MC states are DSD-free states in general local dephasing noise. ■

One can also build other higher dimensional DSD-free states, which could be locally projected onto entangled MC states by following the techniques for lemma 4. As a short summary, lemma 1 to 5 give a few primary results on DSD-free states in finite time, in terms of the special dephasing noise Eq.(1) and general dephasing noise respectively.

To gain a better understanding of the entanglement dynamics, we further study the properties of the evolved state in the infinite-time limit, which is summarized as the following lemma.

Lemma 6. For any qutrit-qutrit state under the dephasing noise in Eq.(1), the final evolved state could be separable or eternally distillable in the infinite-time limit, but no PPT entangled state can exist as time goes to infinity.

Proof. Let us briefly justify the above statement. Consider an arbitrary qutrit-qutrit state σ under the action of an infinite time of local dephasing, then the final state $\sigma(t)$ can always be written as: $a|00\rangle\langle 00| + |0\rangle\langle 0| \otimes (b|1\rangle\langle 1| + c|1\rangle\langle 2| + c^*|2\rangle\langle 1| + d|2\rangle\langle 2|) + (e|1\rangle\langle 1| + f|1\rangle\langle 2| + f^*|2\rangle\langle 1| + g|2\rangle\langle 2|) \otimes |0\rangle\langle 0| + (1|1\rangle\langle 1| + |2\rangle\langle 2|) \otimes (1|1\rangle\langle 1| + |2\rangle\langle 2|) \rho(0) + (1|1\rangle\langle 1| + |2\rangle\langle 2|) \otimes (1|1\rangle\langle 1| + |2\rangle\langle 2|)$. There are four terms in all, and each of them corresponds to one of the terms in Eq.(1), respectively. Note that the first three terms could be removed by the local operator $(|1\rangle\langle 1| + |2\rangle\langle 2|) \otimes (|1\rangle\langle 1| + |2\rangle\langle 2|)$ and it does not change the fourth term. So the evolved state $\sigma(t)$ is entangled if and only if the fourth term is entangled. It immediately implies that the state $\sigma(t)$ could be separable or eternally NPT and distillable in the infinite time limit. As we have seen in the previous sections, both cases also exist in finite time's evolution. Since the entanglement property of the evolved state is determined by the two-qubit state, then we conclude that any PPT entangled state is unstable as time goes to infinity under the local dephasing noise in Eq.(1). In contrast, it's still an open question whether there always exists PPT entangled state in finite time's evolution. ■

IV. DISCUSSIONS AND CONCLUSIONS

Before concluding we would like to discuss some open questions as follow:

Firstly, the initial state in our scheme is a free entangled state. We have shown that it may evolve into a PPT entangled state and then a separable state for finite time; it may also be eternally free entangled for all finite time in the local dephasing noise described by Eq.(1). We have also shown that any qutrit-qutrit PPT entangled state

will lose its entanglement as time goes to infinity in our decoherence model. Then an open question is whether PPT entanglement can be preserved in any finite time under (general) local dephasing noise? If the answer is no (supported by the calculation in Sec.II), then we have another interesting difference between the free and bound entangled state: that is, the former can be robust against the decoherence in any time while the latter cannot.

Secondly, we have proposed a few systematic methods to build DSD-free subspaces for $3 \otimes 3$ state, in which all states keep their distillability asymptotically in local dephasing noise. The primary extension to higher dimensional bipartite system has also been proposed. It's then an open question to find out more entangled states which are robust against different local noises (beyond our model) in any dimensional space. Such states are applicable to quantum information schemes and thus deserve a deeper study.

In summary, we have introduced the concept of DSD through an explicit qutrit-qutrit state. We found that under the bi-local dephasing noise, the free entangled state may evolve into a PPT entangled state (DSD) and then become separable (ESD) permanently. Moreover, a simple local unitary operation on the initial state can avoid

the sudden death of distillability. We also have compared the action of DSD and DSD-free states in terms of the Bures fidelity. Next, we have proposed the systematic methods of building the DSD-free space against decoherence, so that the evolved states are distillable in finite time. Finally, we proved that there is no PPT entangled state in infinite-time limit in our model. Our results imply that further study on free entanglement with time evolution in practical local noises is required.

V. ACKNOWLEDGEMENTS

We thank A. Winter and C. Beny for critical reading the manuscript. W. Song was supported by the Doctoral Startup Foundation of Guangdong Province. The Centre for Quantum Technologies is funded by the Singapore Ministry of Education and the National Research Foundation as part of the Research Centres of Excellence programme. SLZ was supported by NSFC under Grant No. 10674049 and the State Key Program for Basic Research of China(Nos.2006CB921801 and 2007CB925204).

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, New York, 2000).
 - [2] T. Yu and J. H. Eberly, Phys. Rev. Lett. 93, 140404 (2004); *ibid.*, 97, 140403 (2006); Science 323, 598 (2009).
 - [3] M. Yonac *et al.*, J. Phys. B 39, S621 (2006).
 - [4] H. T. Cui *et al.*, Phys. Lett. A 365, 44 (2007).
 - [5] T. Yu and J. H. Eberly, Quantum. Inf. Comput. 7, 459 (2007).
 - [6] F. Lastra *et al.*, Phys. Rev. A 75, 062324 (2007).
 - [7] G.-F. Zhang *et al.*, Opt. Commun. 275, 274 (2007).
 - [8] A. Vaglica and G. Vetri, Phys. Rev. A 75, 062120 (2007).
 - [9] B. Bellomo, R. Lo Franco, and G. Compagno, Phys. Rev. Lett. 99, 160502 (2007); Phys. Rev. A 77, 032342 (2008); J. Dajka, M. Mierzejewski, and J. Luczka, *ibid.* 77, 042316 (2008); X. Cao and H. Zheng, *ibid.* 77, 022320 (2008).
 - [10] M. Ikram, F. L. Li, and M. S. Zubairy, Phys. Rev. A 75, 062336 (2007).
 - [11] A. Al-Qasimi and D. F. V. James, Phys. Rev. A 77, 012117 (2008).
 - [12] Z. Ficek and R. Tanas, Phys. Rev. A 74, 024304 (2006).
 - [13] I. Sainz and G. Bjork, Phys. Rev. A 77, 052307 (2008).
 - [14] K. Ann and G. Jaeger, Phys. Rev. A 76, 044101 (2007); K. Ann and G. Jaeger, Phys. Lett. A 372, 579 (2008).
 - [15] A. R. P. Rau, M. Ali, and G. Alber, Europhys. Lett. 82, 40002 (2008); M. Ali, G. Alber, and A. R. P. Rau, J. Phys. B 42, 025501 (2009).
 - [16] J. H. Huang and S. Y. Zhu, Phys. Rev. A 76, 062322 (2007).
 - [17] P. Li, Q. Zhang, and J. Q. You, Phys. Rev. A 79, 014303 (2009).
 - [18] C.-J. Shan *et al.*, Chin. Phys. Lett. 25, 3115 (2008).
 - [19] Z.-G. Li, *et al.*, Phys. Rev. A 79, 024303 (2009).
 - [20] A. R. R. Carvalho *et al.*, Eur. Phys. J. D 41, 425 (2007).
 - [21] K. Ann and G. Jaeger, Phys. Rev. B 75, 115307 (2007).
 - [22] I. Sainz and G. Bjork, Phys. Rev. A 76, 042313 (2007).
 - [23] C. E. Lopez *et al.*, Phys. Rev. Lett. 101, 080503 (2008).
 - [24] L. Aolita *et al.*, Phys. Rev. Lett. 100, 080501 (2008).
 - [25] Z.-X. Man, Y.-J. Xia, and N. Ba An, Phys. Rev. A 78, 064301 (2008).
 - [26] N. Ba An and J. Kim, Phys. Rev. A 79, 022303 (2009).
 - [27] Z. Liu and H. Fan, Phys. Rev. A 79, 064305 (2009).
 - [28] M. O. Terra Cunha, New J. Phys. 9, 237 (2007).
 - [29] M. P. Almeida *et al.*, Science 316, 579 (2007); A. Salles *et al.*, Phys. Rev. A 78, 022322 (2008).
 - [30] J. Laurat *et al.*, Phys. Rev. Lett. 99, 180504 (2007).
 - [31] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 80, 5239 (1998).
 - [32] D. Yang *et al.*, Phys. Rev. Lett. 95, 190501 (2005).
 - [33] G. Toth *et al.*, Phys. Rev. Lett. 99, 250405 (2007); D. Cavalcanti *et al.*, Phys. Rev. A 78, 012335 (2008).
 - [34] In fact, there are strong evidences showing the existence of bound entangled states with non-positive partial transposition (NPT), see W. Dur *et al.*, Phys. Rev. A 61, 062313 (2000); D. P. DiVincenzo *et al.*, *ibid.*, 61, 062312 (2000). Here, we only consider the PPT bound entangled states.
 - [35] O. Rudolph, Quantum Inf. Process. 4, 219 (2005); K. Chen and L.A. Wu, Quantum. Inf. Comput. 3, 193 (2003).
 - [36] K. Kraus, *State, effects and operations*(Springer-Verlag, Berlin, 1983).
 - [37] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 78, 574 (1997).
 - [38] P. Horodecki, M. Horodecki, and R. Horodecki, Phys.

- Rev. Lett. 82, 1056 (1999).
- [39] D. Bures, Trans. Am. Math. Soc. 135, 199 (1969); A. Uhlmann, Rep. Math. Phys 9, 273 (1976); A. Uhlmann, Rep. Math. Phys. 24, 229 (1986).
- [40] P. Horodecki *et al.*, quant-ph/9910122.
- [41] L. Chen and Y.-X. Chen, Phys. Rev. A 78, 022318 (2008).
- [42] M. Horodecki, A. Sen(De), and U. Sen, Phys. Rev. A 67, 062314 (2003).